



BRIGHT FUTURES HANDOUT ► PARENT

12 MONTH VISIT

Here are some suggestions from Bright Futures experts that may be of value to your family.



HOW YOUR FAMILY IS DOING

- If you are worried about your living or food situation, reach out for help. Community agencies and programs such as WIC and SNAP can provide information and assistance.
- Don't smoke or use e-cigarettes. Keep your home and car smoke-free. Tobacco-free spaces keep children healthy.
- Don't use alcohol or drugs.
- Make sure everyone who cares for your child offers healthy foods, avoids sweets, provides time for active play, and uses the same rules for discipline that you do.
- Make sure the places your child stays are safe.
- Think about joining a toddler playgroup or taking a parenting class.
- Take time for yourself and your partner.
- Keep in contact with family and friends.



FEEDING YOUR CHILD

- Offer healthy foods for meals and snacks. Give 3 meals and 2 to 3 snacks spaced evenly over the day.
- Avoid small, hard foods that can cause choking—popcorn, hot dogs, grapes, nuts, and hard, raw vegetables.
- Have your child eat with the rest of the family during mealtime.
- Encourage your child to feed herself.
- Use a small plate and cup for eating and drinking.
- Be patient with your child as she learns to eat without help.
- Let your child decide what and how much to eat. End her meal when she stops eating.
- Make sure caregivers follow the same ideas and routines for meals that you do.



ESTABLISHING ROUTINES

- Praise your child when he does what you ask him to do.
- Use short and simple rules for your child.
- Try not to hit, spank, or yell at your child.
- Use short time-outs when your child isn't following directions.
- Distract your child with something he likes when he starts to get upset.
- Play with and read to your child often.
- Your child should have at least one nap a day.
- Make the hour before bedtime loving and calm, with reading, singing, and a favorite toy.
- Avoid letting your child watch TV or play on a tablet or smartphone.
- Consider making a family media plan. It helps you make rules for media use and balance screen time with other activities, including exercise.



FINDING A DENTIST

- Take your child for a first dental visit as soon as her first tooth erupts or by 12 months of age.
- Brush your child's teeth twice a day with a soft toothbrush. Use a small smear of fluoride toothpaste (no more than a grain of rice).
- If you are still using a bottle, offer only water.

Helpful Resources: Smoking Quit Line: 800-784-8669 | Family Media Use Plan: www.healthychildren.org/MediaUsePlan

Poison Help Line: 800-222-1222 | Information About Car Safety Seats: www.safercar.gov/parents | Toll-free Auto Safety Hotline: 888-327-4236

12 MONTH VISIT—PARENT

SAFETY

- Make sure your child's car safety seat is rear facing until he reaches the highest weight or height allowed by the car safety seat's manufacturer. In most cases, this will be well past the second birthday.
- Never put your child in the front seat of a vehicle that has a passenger airbag. The back seat is safest.
- Place gates at the top and bottom of stairs. Install operable window guards on windows at the second story and higher. Operable means that, in an emergency, an adult can open the window.
- Keep furniture away from windows.
- Make sure TVs, furniture, and other heavy items are secure so your child can't pull them over.
- Keep your child within arm's reach when he is near or in water.
- Empty buckets, pools, and tubs when you are finished using them.
- Never leave young brothers or sisters in charge of your child.
- When you go out, put a hat on your child, have him wear sun protection clothing, and apply sunscreen with SPF of 15 or higher on his exposed skin. Limit time outside when the sun is strongest (11:00 am–3:00 pm).
- Keep your child away when your pet is eating. Be close by when he plays with your pet.
- Keep poisons, medicines, and cleaning supplies in locked cabinets and out of your child's sight and reach.
- Keep cords, latex balloons, plastic bags, and small objects, such as marbles and batteries, away from your child. Cover all electrical outlets.
- Put the Poison Help number into all phones, including cell phones. Call if you are worried your child has swallowed something harmful. Do not make your child vomit.

WHAT TO EXPECT AT YOUR CHILD'S 15 MONTH VISIT

We will talk about

- Supporting your child's speech and independence and making time for yourself
- Developing good bedtime routines
- Handling tantrums and discipline
- Caring for your child's teeth
- Keeping your child safe at home and in the car

490 Hwy 85N, Ste A
Niceville, FL 32578
(850) 424-6208

Beachside Pediatric
Associates, P.A.



Consistent with *Bright Futures: Guidelines for Health Supervision of Infants, Children, and Adolescents, 4th Edition*

For more information, go to <https://brightfutures.aap.org>.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



The information contained in this handout should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances. Original handout included as part of the *Bright Futures Tool and Resource Kit, 2nd Edition*.

Inclusion in this handout does not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of the resources mentioned in this handout. Web site addresses are as current as possible but may change at any time.

The American Academy of Pediatrics (AAP) does not review or endorse any modifications made to this handout and in no event shall the AAP be liable for any such changes.

© 2019 American Academy of Pediatrics. All rights reserved.

Immunizations: What You Need to Know

Vaccines (immunizations) keep children healthy. Vaccines are safe. Vaccines are effective. Vaccines save lives.

However, parents may still have questions about why vaccines are needed, and some parents may be concerned about vaccine safety because they have been misinformed.

Read on for answers from the American Academy of Pediatrics (AAP) to some common questions parents have about vaccines. The AAP is a source you can trust for reliable medical information.

Q: What vaccines does my child need?

A: Children need all the following vaccines to stay healthy:

- **Hepatitis A and hepatitis B vaccines** to help protect against serious liver diseases.
- **Rotavirus vaccine** to help protect against the most common cause of diarrhea and vomiting in infants and young children. Rotavirus is the most common cause of hospitalizations in young infants due to vomiting, diarrhea, and dehydration.
- **DTaP and Tdap vaccines** to help protect against diphtheria, tetanus (lockjaw), and pertussis (whooping cough).
- **Hib vaccine** to help protect against *Haemophilus influenzae* type b (a cause of spinal meningitis and other serious infections).
- **Pneumococcal vaccine** to help protect against bacterial meningitis, pneumonia, and infections of the blood.
- **Polio vaccine** to help protect against a crippling viral disease that can cause paralysis.
- **Influenza vaccine** to help protect against influenza (flu), a potentially fatal disease. This vaccine is recommended for all people beginning at 6 months and older.
- **MMR vaccine** to help protect against measles, mumps, and rubella (German measles), all highly contagious and potentially very serious diseases.
- **Varicella vaccine** to help protect against chickenpox and its many complications, including flesh-eating strep, staph toxic shock, and encephalitis (an inflammation of the brain).
- **Meningococcal vaccine** to help protect against very serious bacterial diseases that affect the blood, brain, and spinal cord.
- **HPV (human papillomavirus) vaccine** to prevent cancers of the mouth and throat, cervix, and genitals.

Remember, vaccines prevent diseases and save lives. It's important to follow the schedule recommended by the AAP. Contact your child's doctor if you have any questions.

Q: Why are some of these vaccines still needed if the diseases are not as common anymore?

A: Many of these diseases are not as common as they once were because of vaccines. However, the bacteria and viruses that cause them still exist and can still make children very sick.

For example, before the Hib vaccine was developed in the 1980s, there were about 20,000 cases of Hib disease in the United States a year. Today there are fewer than 100 cases a year. However, the bacteria that causes Hib disease still exists. That is why children still need the vaccine to be protected.

In the United States, vaccines protect children from many diseases. However, in many parts of the world vaccine-preventable diseases are still common. Because diseases may be brought into the United States by Americans who travel abroad or by people visiting areas with current disease outbreaks, it's important that your child is vaccinated.

Q: Chickenpox is not a fatal disease, so why is the vaccine needed?

A: Chickenpox is usually mild. However, there can be serious complications. In fact, before the vaccine was licensed in 1995, there were about 4 million cases, 11,000 hospitalizations, and 100 deaths each year from chickenpox. Chickenpox is also very contagious. Most children feel miserable and miss 1 week or more of school when infected. It is because of the vaccine that the number of cases of chickenpox and its complications, including deaths, have gone down so dramatically.

Q: Does my baby need immunizations if I am breastfeeding?

A: Yes. While breastfeeding gives some protection against many diseases (and is the best nutrition for your baby), it is not a substitute for vaccines. In fact, breastfeeding and vaccines work well together. Studies show that breastfed babies respond better to vaccines and get better protection from them than babies who are not breastfed. And breastfeeding during or right after immunizations may help calm babies upset by the shots.


Q: Do vaccines even work? It seems like most of the people who get these diseases have been vaccinated.

A: Yes. Vaccines work very well. Millions of children have been protected against serious illnesses because they were immunized. Most childhood vaccines are 90% to 99% effective in preventing disease. Children who aren't vaccinated are much more likely to get a disease if they are exposed to it. And if a vaccinated child does get the disease, the symptoms are usually milder with fewer complications than in a child who hasn't been vaccinated.

Q: When should my child get immunized?

A: Children should get most of their shots during their first 2 years after birth. This is because many of these diseases are the most severe in the very young. Most newborns receive their first shot (hepatitis B) at birth before leaving the hospital, and more are given at well-child checkups in the first 6 months after birth. Other shots are given before children go to school. Older children and teens need vaccines to continue to protect them throughout adolescence and early adulthood. (Parents and caregivers also need vaccines so that they can prevent bringing infections home to their children and to keep themselves healthy so that they can care for their children!)

Children who are not immunized or who are behind on their shots are at risk of getting many of these diseases. They can also spread these diseases to others who have not yet been immunized. Ask your child's doctor if your child is up to date. Keep track of the vaccines each child receives and bring this information to each doctor visit.



Q: What side effects will my child have after getting a vaccine? Are they serious?

A: There may be mild side effects, such as swelling, redness, and tenderness where the shot was given, but they do not last long. Your child may also have a slight fever and be fussy for a short time afterward. Your doctor may suggest giving your child pain medicine to help relieve discomfort. It is very rare for side effects to be serious. However, you should call your child's doctor if you have any concerns after vaccines are given.

Q: Should some children not be immunized?

A: Children with certain health problems may need to avoid some vaccines or get them later. In most cases, children with cancer, those taking oral or injected steroids for lung or kidney conditions, or those who have problems with their immune systems should not get vaccines that are made with live viruses. To protect these children, it is very important for others to be vaccinated. On the other hand, a child with a minor illness, such as low-grade fever, an ear infection, cough, a runny nose, or mild diarrhea, can safely be immunized.

Q: Does the MMR vaccine cause autism?

A: No! The MMR vaccine does not cause autism spectrum disorder (ASD). Many research studies have been done to address this issue. There may be confusion because children with ASD are often diagnosed between 18 and 30 months of age—around the same time the MMR vaccine is given. This has led some people to assume that the vaccine is the cause. Increasing evidence shows that even though the symptoms of ASD may not be visible until the second year after birth or later, ASD starts before a baby is born.

Q: Do vaccines cause SIDS?

A: No! Babies get many of their first vaccines between 2 and 4 months of age. This is also the peak age for sudden infant death syndrome (SIDS), which is why some people feel they might be related. However, careful scientific studies have confirmed that vaccinations not only do not cause SIDS but may help prevent it.

Q: How do we know vaccines are safe?

A: The safety and effectiveness of vaccines are under constant study. Because vaccines are designed to be given routinely during well-child visits, they must be safe. Safety testing begins as soon as a new vaccine is considered, continues until it is approved by the US Food and Drug Administration (FDA), and is monitored indefinitely after licensure. The AAP works closely with the Centers for Disease Control and Prevention (CDC) to make recommendations for vaccine use.

Q: What is thimerosal and does it cause neurologic problems?

A: In the 1930s a preservative called thimerosal was added to vaccines to prevent contamination of vaccines. Thimerosal contains very small amounts of mercury, but it is in a different form than the potentially harmful mercury we are all exposed to in the environment. Even after many studies, the type of mercury in thimerosal has never been shown to cause health problems other than rare allergic reactions in some people. Thimerosal does not cause neurologic problems. Since 2001 all vaccines for infants either are thimerosal-free or contain only trace amounts of the preservative. Many are available in single-dose, preservative-free forms.

Q: Is it safe to give more than one vaccine at a time?

A: Yes! Your child's immune system is capable of handling multiple vaccines. Many years of experience and careful research have shown that routine childhood vaccines can be given together safely and effectively. Side effects are not increased when vaccines are given together.

Q: Where can I find more information?

A: Be sure your information comes from reliable and accurate sources. You cannot trust everything you find on the internet. Credible sources include

American Academy of Pediatrics

www.aap.org and www.HealthyChildren.org

CDC Vaccines & Immunizations

www.cdc.gov/vaccines

Immunization Action Coalition

www.immunize.org

Remember

If you have any questions or concerns about your child's health, contact your child's doctor.

American Academy of Pediatrics
DEDICATED TO THE HEALTH OF ALL CHILDREN®



The American Academy of Pediatrics (AAP) is an organization of 67,000 primary care pediatricians, pediatric medical subspecialists, and pediatric surgical specialists dedicated to the health, safety, and well-being of infants, children, adolescents, and young adults.

Any websites, brand names, products, or manufacturers are mentioned for informational and identification purposes only and do not imply an endorsement by the American Academy of Pediatrics (AAP). The AAP is not responsible for the content of external resources. Information was current at the time of publication. The information contained in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.

© 2020 American Academy of Pediatrics. All rights reserved.

Immunizations for Babies

A Guide for Parents

These are the vaccinations your baby needs!

| | |
|---------------------|--|
| At birth | HepB |
| 2 months | HepB ¹ + DTaP + PCV13 + Hib + Polio + RV |
| 4 months | HepB ² + DTaP + PCV13 + Hib + Polio + RV |
| 6 months | HepB ¹ + DTaP + PCV13 + Hib ³ + Polio + RV ⁴ + Influenza ⁵ |
| 12 months and older | MMR + DTaP + PCV13 + Hib + Chickenpox + HepA ⁶ + Influenza ⁵ |

Check with your doctor or nurse to make sure your baby is receiving all vaccinations on schedule. Many times vaccines are combined to reduce the number of injections. Be sure you ask for a record card with the dates of your baby's vaccinations; bring this with you to every visit.

Here's a list of the diseases your baby will be protected against:

HepB: hepatitis B, a serious liver disease

DTaP: diphtheria, tetanus (lockjaw), and pertussis (whooping cough)

PCV13: pneumococcal conjugate vaccine protects against a serious blood, lung, and brain infection

Hib: *Haemophilus influenzae* type b, a serious brain, throat, and blood infection

Polio: polio, a serious paralyzing disease

RV: rotavirus infection, a serious diarrheal disease

Influenza: a serious lung infection

MMR: measles, mumps, and rubella

HepA: hepatitis A, a serious liver disease

Chickenpox: also called varicella


 490 Hwy 85N, Ste A
 Niceville, FL 32578
 (850) 424-6208
 Beachside Pediatric
 Associates, P.A.



Notes to above chart:

1. This is the age range in which this vaccine should be given.
2. Your baby may not need a dose of Hep B vaccine at age 4 months, depending on the vaccine used. Check with your doctor or nurse.
3. Your baby may not need a dose of Hib vaccine at age 6 months, depending on the vaccine used. Check with your doctor or nurse.
4. Your baby may not need a dose of RV vaccine at age 6 months, depending on the vaccine used. Check with your doctor or nurse.
5. All children age 6 months and older should be vaccinated against influenza in the fall or winter of each year.
6. Your child will need 2 doses of HepA vaccine, given at least 6 months apart.

immunization
action coalition



Saint Paul, Minnesota • 651-647-9009 • www.immunize.org • www.vaccineinformation.org

www.immunize.org/catg.d/p4010.pdf • Item #P4010 (8/20)

Table 1

Recommended Child and Adolescent Immunization Schedule for ages 18 years or younger, United States, 2021

These recommendations must be read with the notes that follow. For those who fall behind or start late, provide catch-up vaccination at the earliest opportunity as indicated by the green bars. To determine minimum intervals between doses, see the catch-up schedule (Table 2). School entry and adolescent vaccine age groups are shaded in gray.

| Vaccine | Birth | 1 mo | 2 mos | 4 mos | 6 mos | 9 mos | 12 mos | 15 mos | 18 mos | 19–23 mos | 2–3 yrs | 4–6 yrs | 7–10 yrs | 11–12 yrs | 13–15 yrs | 16 yrs | 17–18 yrs | |
|---|----------------------|--------------------------|----------------------|----------------------|-----------|-----------|--|--------------------------|--------------------------|---------------------------------|---------|---------------------------------|----------|-----------|-----------|--------|--------------------------------|----------------------|
| Hepatitis B (HepB) | 1 st dose | ← 2 nd dose → | | | | | | | ← 3 rd dose → | | | | | | | | | |
| Rotavirus (RV): RV1 (2-dose series), RV5 (3-dose series) | | | 1 st dose | 2 nd dose | See Notes | | | | | | | | | | | | | |
| Diphtheria, tetanus, acellular pertussis (DTaP <7 yrs) | | 1 st dose | 2 nd dose | 3 rd dose | | | | | ← 4 th dose → | | | 5 th dose | | | | | | |
| Haemophilus influenzae type b (Hib) | 1 st dose | 2 nd dose | See Notes | | | | ← 3 rd or 4 th dose, See Notes → | | | | | | | | | | | |
| Pneumococcal conjugate (PCV13) | 1 st dose | 2 nd dose | 3 rd dose | | | | ← 4 th dose → | | | | | | | | | | | |
| Inactivated poliovirus (IPV <18 yrs) | 1 st dose | 2 nd dose | | | | | | | ← 3 rd dose → | | | 4 th dose | | | | | | |
| Influenza (IIV) | | | | | | | | | | Annual vaccination 1 or 2 doses | | | | | | | | |
| or | | | | | | | | | | | | | | | | | | |
| Influenza (LAIV4) | | | | | | | | | | | | Annual vaccination 1 or 2 doses | | | | | Annual vaccination 1 dose only | |
| Measles, mumps, rubella (MMR) | | | | | | See Notes | ← 1 st dose → | | | | | 2 nd dose | | | | | | |
| Varicella (VAR) | | | | | | | ← 1 st dose → | | | | | 2 nd dose | | | | | | |
| Hepatitis A (HepA) | | | | | | See Notes | | 2-dose series, See Notes | | | | | | | | | | |
| Tetanus, diphtheria, acellular pertussis (Tdap ≥7 yrs) | | | | | | | | | | | | | | | | | Tdap | |
| Human papillomavirus (HPV) | | | | | | | | | | | | | | * | See Notes | | | |
| Meningococcal (MenACWY-D ≥9 mos, MenACWY-CRM ≥2 mos, MenACWY-TT ≥2years) | | | | | | | | | | | | | | | | | 1 st dose | 2 nd dose |
| Meningococcal B | | | | | | | | | | | | | | | | | | See Notes |
| Pneumococcal polysaccharide (PPSV23) | | | | | | | | | | | | | | | | | | See Notes |

Range of recommended ages for all children

Range of recommended ages for catch-up immunization

Range of recommended ages for certain high-risk groups

Recommended based on shared clinical decision-making or *see footnote 1

No recommendation/ not applicable

After the Shots...

Your child may need extra love and care after getting vaccinated. Some vaccinations that protect children from serious diseases also can cause discomfort for a while. Here are answers to questions many parents have after their children have been vaccinated. If this sheet doesn't answer your questions, call your healthcare provider.

Vaccinations may hurt a little... but disease can hurt a lot!

Call your healthcare provider right away if you answer "yes" to any of the following questions:

- Does your child have a temperature that your healthcare provider has told you to be concerned about?
- Is your child pale or limp?
- Has your child been crying for more than 3 hours and just won't quit?
- Is your child's body shaking, twitching, or jerking?
- Is your child very noticeably less active or responsive?

► Please see page 2 for information on the proper amount of medicine to give your child to reduce pain or fever.

What to do if your child has discomfort

I think my child has a fever. What should I do?

Check your child's temperature to find out if there is a fever. An easy way to do this is by taking a temperature in the armpit using an electronic thermometer (or by using the method of temperature-taking your healthcare provider recommends). If your child has a temperature that your healthcare provider has told you to be concerned about or if you have questions, call your healthcare provider.

Here are some things you can do to help reduce fever:

- Give your child plenty to drink.
- Dress your child lightly. Do not cover or wrap your child tightly.
- Give your child a fever- or pain-reducing medicine such as acetaminophen (e.g., Tylenol) or ibuprofen (e.g., Advil, Motrin). The dose you give your child should be based on your child's weight and your healthcare provider's instructions. See the dose chart on page 2. *Do not give aspirin.* Recheck your child's temperature after 1 hour. Call your healthcare provider if you have questions.

My child has been fussy since getting vaccinated. What should I do?

After vaccination, children may be fussy because of pain or fever. To reduce discomfort, you may want to give your child a medicine such as acetaminophen or ibuprofen. See the dose chart on page 2. *Do not give aspirin.* If your child is fussy for more than 24 hours, call your healthcare provider.

My child's leg or arm is swollen, hot, and red. What should I do?

- Apply a clean, cool, wet washcloth over the sore area for comfort.
- For pain, give a medicine such as acetaminophen or ibuprofen. See the dose chart on page 2. *Do not give aspirin.*
- If the redness or tenderness increases after 24 hours, call your healthcare provider.

My child seems really sick. Should I call my healthcare provider?

If you are worried **at all** about how your child looks or feels, call your healthcare provider!

HEALTHCARE PROVIDER: PLEASE FILL IN THE INFORMATION BELOW.

If your child's temperature is 102.5°F or 39°C or higher, or if you have questions, call your healthcare provider.

Healthcare provider phone number (850) 424-6208

immunization
action coalition

A

immunize.org

Saint Paul, Minnesota • 651-647-9009 • www.immunize.org • www.vaccineinformation.org

www.immunize.org/catg.d/p4015.pdf • Item #P4015 (2/19)

Start Reading to Your Child Early



How to Help Your Child Learn to Read

A baby can enjoy books by 6 months of age! Here are things you can do with your child at different ages to help your child learn to love words and books.

Birth to Age 1

- Play with your baby often. Talk, sing, and say rhymes. This helps your baby learn to talk.
- Talk with your baby, making eye contact. Give your baby time to answer in baby talk.
- Give your baby sturdy board books to look at. It's OK for a baby to chew on a book.
- Look at picture books with your baby and name things. Say "See the baby!" or "Look at the puppy!"
- Babies like board books with pictures of babies and everyday objects like balls and blocks.
- Snuggle with your baby on your lap and read aloud. Your baby may not understand the story, but will love the sound of your voice and being close to you.
- Don't let your child watch TV until age 2 or older.

1 to 3 Years of Age

- Read to your child every day. Let your child pick the book, even if it's the same one again and again!
- Younger toddlers (1 to 2 years of age) like board books with pictures of children doing everyday things (like eating and playing). They also like "goodnight" books and books with rhymes. Books should only have a few words on each page.

- Older toddlers (2 to 3 years of age) like board books and books with paper pages. They love books with rhymes and words that are repeated. Books about families, friends, animals, and trucks are also good.
- Let your child "read" to you by naming things in the book or making up a story.
- Take your child to the library. Celebrate your child getting a library card!
- Keep talking, singing, saying rhymes, and playing with your child.
- Don't let your child watch TV until age 2 or older.

Reading Tips

- Set aside time every day to read together. Reading at bedtime is a great way to get ready for sleep.
- Leave books in your children's rooms for them to enjoy on their own. Have a comfortable bed or chair, bookshelf, and reading lamp.
- Read books your child enjoys. Your child may learn the words to a favorite book. Then, let your child complete the sentences, or take turns saying the words.
- Don't drill your child on letters, numbers, colors, shapes, or words. Instead, make a game of it.

490 Hwy 85N, Ste A
 Niceville, FL 32578
 (850) 424-6208

Beachside Pediatric Associates, P.A.



Continued on back

Continued from front

3 to 5 Years of Age

- Read ABC books with your child. Point out letters as you read.
- Preschool children like books that tell stories. They also love counting books, alphabet books, and word books. Like toddlers, they love books with rhymes and words they can learn by heart.
- Help your child recognize whole words as well as letters. Point out things like letters on a stop sign or the name on a favorite store.
- Ask your child questions about the pictures and story. Invite him or her to make up a story about what's in the book.
- Some public TV shows, videos, and computer games can help your child learn to read. But you need to be involved too. Watch or play *with* your child and talk about the program. Limit TV time to 1 or 2 hours per day. Avoid violent shows and movies. Try to stick to educational shows.
- Give your child lots of chances to use written words. Write shopping lists together. Write letters to friends or family.



Read Aloud With Your Child

Reading aloud is one of the best ways to help your child learn to read. The more excited you act when you read a book, the more your child will enjoy it.

- Use funny voices and animal noises!
- Look at the pictures. Ask your child to name things in the pictures. Talk about how the pictures go with the story. Ask what is happening in the story.

- Invite your child to join in when a line is repeated over and over.
- Show your child how things in the book are like things in your child's life.
- If your child asks a question, stop and answer it. Books can help children express their thoughts and solve problems.
- Keep reading to your child even after he or she learns to read. Children can listen and understand harder stories than they can read on their own.

Listen to Your Child Read Aloud

Once your child starts reading, have him or her read out loud. Take turns reading.

If your child asks for help with a word, give it right away. But let your child sound out words if he or she wants to.

Know when your child has had enough. Stop if your child is tired or frustrated.

Most of all, give lots of praise! You are your child's first, and most important, teacher!

The American Academy of Pediatrics (AAP) is grateful for the Reach Out and Read program's help with this handout. Reach Out and Read works with children's doctors to make promoting literacy and giving out books part of children's basic health care. This program is endorsed by the AAP. To learn more about Reach Out and Read, go to www.reachoutandread.org.

181 West 12th St A
 Merrill, FL 32579
 (352) 426-4108





To learn more, visit the American Academy of Pediatrics (AAP) Web site at www.aap.org.
 Your child's doctor will tell you to do what's best for your child. This information should not take the place of talking with your child's doctor.
 We hope the resources in this handout are helpful. The AAP is not responsible for the information in these resources. We try to keep the information up to date but it may change at any time.
 Adaptation of the AAP information in this handout into plain language was supported in part by McNeil Consumer Healthcare.
 © 2008 American Academy of Pediatrics

American Academy
 of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

1 to 2 Years



1/2



1 TO 2 YEARS

Safety for Your Child

Did you know that injuries are the leading cause of death of children in the United States? Most of these injuries can be prevented.

Often, injuries happen because parents are not aware of what their children can do. At this age your child can *walk, run, climb, jump, and explore* everything. Because of all the new things he or she can do, this stage is a very dangerous time in your child's life. It is your responsibility to protect your child from injury. Your child cannot understand danger or remember "no" while exploring.

Firearm Hazards

Children in homes where guns are present are in more danger of being shot by themselves, their friends, or family members than of being injured by an intruder. It is best to keep all guns out of the home. **Handguns are especially dangerous.** If you keep a gun, keep it unloaded and in a locked place, with the ammunition locked separately. Ask if the homes where your child visits or is cared for have guns and how they are stored.



Poisonings

Children continue to explore their world by putting everything in their mouths, even if it doesn't taste good. Your child can *open doors and drawers, take things apart, and open bottles* easily now, so you must use safety caps on all medicines and toxic household products. **Keep the safety caps on** at all times or find safer substitutes to use. Contact Poison Help for more information.




Your child is now able to get into and on top of everything. Be sure to keep all household products and medicines completely out of sight and reach. Never store lye drain cleaners in your home. Keep all products in their original containers. Use medicines exactly as directed and dispose of unused medicine safely as soon as you are finished with it.

If your child does put something poisonous into his or her mouth, call Poison Help immediately. Add the Poison Help line (1-800-222-1222) to your phone contacts list. Do not make your child vomit.

Falls

To prevent serious falls, lock the doors to any dangerous areas. **Use gates on stairways and install operable window guards** above the first floor. **Remove sharp-edged furniture** from the room your child plays and sleeps in. At this age your child will walk well and start to climb, jump, and run as well. A chair left next to a kitchen counter, table, or window allows your child to climb to dangerously high places. Remember, your child does not understand what is dangerous.



If your child has a serious fall or does not act normally after a fall, call your doctor.

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN®



(over)

Burns

The kitchen is a dangerous place for your child during meal preparation. Hot liquids, grease, and hot foods spilled on your child will cause serious burns. A **safer place for your child** while you are cooking, eating, or unable to give him or her your full attention is the **playpen, crib, or stationary activity center, or buckled into a high chair**. It's best to keep your child out of the kitchen while cooking.



Children who are learning to walk will grab anything to steady themselves, including hot oven doors, wall heaters, or outdoor grills. Keep your child out of rooms where there are hot objects that may be touched, or put a barrier around them. If you have a gas fireplace, keep children away while it is in use and for at least an hour after turning it off. The glass doors get extremely hot and can cause severe burns.

Your child will *reach* for your hot food or cup of coffee, so don't leave it within your child's reach. **NEVER carry your child and hot liquids at the same time.** You can't handle both.

If your child does get burned, immediately put cold water on the burned area. Keep the burned area in cold water for a few minutes to cool it off. Then cover the burn loosely with a dry bandage or clean cloth. Call your doctor for all burns. To protect your child from tap water scalds, the hottest temperature at the faucet should be no more than 120°F. In many cases you can adjust your water heater.

Make sure you have a working smoke alarm on every level of your home, especially in furnace and sleeping areas. Test the alarms every month. It is best to use smoke alarms that use long-life batteries, but if you do not, change the batteries at least once a year.

Drowning


At this age your child loves to play in water. **NEVER leave your child alone in or near a bathtub, pail of water, wading or swimming pool, or any other water, even for a moment.** Empty all buckets after each use. Keep the bathroom doors closed. Your child can drown in less than 2 inches of water. Knowing how to swim does NOT mean your child is safe near or in water. Stay within an arm's length of your child around water.



If you have a swimming pool, fence it on all 4 sides with a fence at least 4 feet high, and be sure the gates are self-latching. If possible, lock doors that could lead to the pool area. Most children drown when they wander out of the house and fall into a pool that is not fenced off from the house. You cannot watch your child every minute while he or she is in the house. It only takes a moment for your child to get out of your house and fall into your pool.

And Remember Car Safety

Car crashes are a great danger to your child's life and health. The crushing forces to your child's brain and body in a crash or sudden stop, even at low speeds, can cause severe injuries or death. **To prevent these injuries USE a car safety seat EVERY TIME** your child rides in the car. All infants and toddlers should ride in a rear-facing car safety seat until they reach the highest weight or height allowed by their car safety seat's manufacturer. Be sure that the safety seat is installed and used correctly. Read and follow the instructions that come with the car safety seat and the instructions for using car safety seats in the owners' manual of your car. **The safest place for all infants and children to ride is in the back seat.**



Do not leave your child alone in or around the car. Keep vehicles and their trunks locked. Children who are left in a car can die of heatstroke because temperatures can reach deadly levels in minutes. They can be strangled by power windows or knock the vehicle into gear.

Always **walk behind your car** to be sure your child is not there before you back out of your driveway. You may not see your child behind your car in the rearview mirror.

From Your Doctor

Remember, the biggest threat to your child's life and health is an injury.

The information in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on individual facts and circumstances.